

# Study of Sleep Quality and its Relation with Educational Status of Iran University of Medical Sciences Students

Seyedeh Melika Kharghani Moghaddam<sup>1</sup>, Iraj Alimohammadi<sup>2</sup>, Shahram Vosoughi<sup>2</sup>, Seyed Hosein Tabatabaei<sup>2</sup>, Rana Ghasemi<sup>2</sup>, Mahdi Ahmadian<sup>2</sup>, Hossein Ebrahimi<sup>3\*</sup>

<sup>1</sup>Department of health education, school of public health, Iran University of Medical Sciences, Tehran, Iran. <sup>2</sup>Department of occupational health, school of public health, Iran University of Medical Sciences, Tehran, Iran. <sup>3</sup>Air pollution research center, Iran University of Medical Sciences, Tehran, Iran. \*corresponding author: Hossein Ebrahimi, Email:hossein.ebrahimi@yahoo.com Tel:+98-912-5117749

## Abstract

**Background:** Undoubtedly, lack of good quality sleep is a major problem for students, affecting their educational success causing them academic failure. Therefore, this study is carried out aiming to determine sleep quality and its relation with educational status of Iran University of Medical Sciences students. **Methods:** This study is a sectional-analytic one performed on dormitory and non-dormitory students of Iran University of Medical Sciences. The criteria of entering the study include being student for at least one year, non-affliction to special diseases and not using drugs and tobacco. People were selected in proportional stratified sampling method after determining the sample volume (400), and Petersburg sleep quality questionnaires were distributed among them. After completion of the questionnaires, their data were extracted and introduced into SPSS version 21. The data obtained via T statistical tests for which Pearson correlation was analyzed were independent. **Results:** The sleep quality mark was higher than 5 for majority of the students including the dormitory and non-dormitory ones; expressing weak sleep quality. Sleep quality was 7.36 for dormitory students and 7.97 for non-dormitory ones; expressing significant statistical difference ( $p=0.018$ ). The results suggest that educational status of non-dormitory students (0.22) is better than the dormitory ones (-1.23), and the difference is statistically significant ( $p=0.000$ ). moreover, the results suggest that there is no relation between good sleep quality and educational status of the students ( $p=0.232$ ). **Conclusion:** The study results suggest that sleep quality is not related to educational status. Considering that educational status is a complex matter and is affected by multiple factors including: intelligence, motivation and hard work, sleep quality plays less significant role regarding educational status.

**Key words:** Educational Status; Sleep Quality; Residence; Dormitory; Non-Dormitory

## Introduction

Desirable sleep is an ensuring factor for mental and physical health which helps with coming into compromise with the others.

This concept helps with increasing cognitive function of people and their focusing in daily activities. Sleep disorder on the other hand causes exhaustion,

**Citation:** Kharghani Moghaddam SM, Alimohammadi I, Vosoughi Sh, Tabatabaei SH, Ghasemi R, Ahmadian M, et al. **Study of Sleep Quality and its Relation with Educational Status of Iran University of Medical Sciences Students.** Archives of Occupational Health. 2019; 3(4): 468-73.

**Article History:** Received: 1 July 2019; Revised: 2 August 2019; Accepted: 10 September 2019

**Copyright:** ©2019 The Author(s); Published by Shahid Sadoughi University of Medical Sciences. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

headache, lack of focus, problem solving ability deterioration, malfunctioning in cognitive functions and life expectancy quality decrement. Additionally, it can diminish problem solving merit or even speech ability. Moreover, ability of learning complex cognitive tasks and skill techniques diminish decrease because of sleep disorder.<sup>1,4</sup> Human health is related to sleep quality and quantity. Night time sleeplessness can affect the life quality, up to such as increasing chance of getting depression and increasing anxiety, as well as weakening ability of encountering daily stresses.<sup>5</sup> In such situations, one becomes angry and provoked easily. Night time sleep quality and quantity can affect the cognitive function and concentration level when doing daily activities.<sup>6</sup>

In addition, morning wakeup time can also affect the daily performance. According to a research performed by Lima et al in Brazil, it was discovered that the group of medical science students who wakeup very early in the morning, do not have good daily performance;<sup>7</sup> while in another researched carried out by Jones et al, it was concluded that performance weakness is related to waking up late in the morning.<sup>8</sup> Hence, it seems there is no clear agreement about waking up time. In the other hand, multiple researches suggest that night time sleep disorder can weaken the daily performance. The Hidalgo et al research in France is of such researches.<sup>9</sup> According to the studies carried out, it is said that sleep disorder complaining among students of medical branches is a common matter those afflicted with rarely seek for therapy.<sup>10</sup> Skipping therapy by the students does not only include sleep disorder; it also covers other psychological disorders as well.<sup>11,12</sup>

Educational success of students is an essential matter in education which can cause decrement of science level of the future students if neglected. Considering that the graduated students of these fields must guarantee health and sanity of the society, significance of this matter is not to be neglected. Undoubtedly, a major problem for the majority of the students affecting their educational success and

causing educational failure is lack of sleep. Educational failure of the youth is not only a personal problem, but is also a social problem to solve which, fundamental measures must be taken. Because problems caused by educational failure of the students will involve the society in the future. This study is carried out aiming to study sleep quality and its relation with educational status of the Iran University of Medical Sciences students.

## Methods

The present study is a sectional-analytic one performed on students of the Iran University of Medical Sciences. Calculation of the sample volume is estimated using sample volume formula for prevalence studies considering the minimum prevalence in other universities in student community at  $P=38\%$  and  $5\%$  error and also assuming accuracy of  $d=0.05$  for minimum sample volume of 399. Statistical society of this study was all students studying in the Iran University of Medical Sciences during the data collection. Therefore, according to the list of students of each faculty and academic grades, 400 of them were selected with proportional stratified sampling method on condition of meeting the required criteria for entering the study which were being student for at least one year, no mental or physical diseases, not using tobacco or drugs and full consent on participating in the study.

The Petersburg sleep quality questionnaire was used to gather data. Several questions were added to the beginning of this questionnaire regarding evaluation of anthropological characteristics (age, gender, marital status, occupation status, university entering year) and educational status. The educational status was evaluated as self-reporting based on a -6 to 6 scale in comparison to the high school time. One of the best tools designed and made regarding sleep quality is the Petersburg sleep quality questionnaire which was formed in 1989 by Dr. Boys et al in Petersburg Psychology institute consisting of 9 items originally, but since question 5 includes 10 peripheral items, the entire questionnaire contains 19

items overall which are to be marked in a 4-degrees Likert scale.<sup>13</sup> This questionnaire includes 7 sub-scales including: subjective sleep quality, sleep latency, sleep duration, habitual sleep efficiency, sleep disturbances, use of sleeping medication and daytime dysfunction. Achieving mark higher than 5 in the entire questionnaire means weak sleep quality.<sup>13</sup>

Boys et al (1989) who made and introduced this questionnaire for the first time, obtained the 0.83 for internal integrity of the questionnaire using Cronbach's alpha.<sup>13</sup> Justifiability and stability were obtained 0.86 and 0.89 respectively in the Iranian version of the questionnaire.<sup>14</sup> After determining the random sample and society, the prepared questionnaire is given to the studied students to fill in self-report method. After filling in the questionnaires, their information was

extracted and introduced into SPSS version 21. After gathering the data, they are analyzed by independent statistical T and Pearson correlation tests. Significance level of tests was considered 0.05.

### Results

Four hundred students of the Iran university of Medical Sciences were selected for this study in random with age average of 23.97(5.95) years old. Results of overall sleep status and its dimensions among the studied students based on residence status is included in table1. In this table, the 7-dimensions sleep quality in 4 classes (no sleep problems, medium sleep problems, serious sleep problem and very serious sleep problem) and overall sleep quality in two classes of weak and desired sleep quality are expressed.

**Table 1.** frequency distribution of overall sleep quality ant its dimensions for the studied students based on residency status

| Dimension                   | Classification              | Residence status |                      |               |                      |
|-----------------------------|-----------------------------|------------------|----------------------|---------------|----------------------|
|                             |                             | Dormitory        |                      | Non-dormitory |                      |
|                             |                             | Number           | Authentic percentage | Number        | Authentic percentage |
| 1 Overall sleep status      | Desired                     | 40               | 20                   | 27            | 14                   |
|                             | Weak                        | 156              | 78                   | 168           | 84                   |
| 2 Subjective sleep quality  | No sleep problems           | 38               | 19                   | 26            | 13                   |
|                             | Medium sleep problems       | 115              | 57                   | 127           | 64                   |
|                             | Serious sleep problems      | 39               | 19                   | 29            | 15                   |
|                             | Very serious sleep problems | 9                | 5                    | 15            | 8                    |
| 3 Sleep latency             | No sleep problems           | 35               | 17                   | 25            | 13                   |
|                             | Medium sleep problems       | 91               | 45                   | 89            | 45                   |
|                             | Serious sleep problems      | 52               | 26                   | 62            | 31                   |
|                             | Very serious sleep problems | 21               | 10                   | 21            | 11                   |
| 4 Sleep duration            | No sleep problems           | 84               | 42                   | 78            | 39                   |
|                             | Medium sleep problems       | 55               | 27                   | 57            | 29                   |
|                             | Serious sleep problems      | 33               | 16                   | 33            | 17                   |
|                             | Very serious sleep problems | 28               | 14                   | 31            | 15.6                 |
| 5 Habitual sleep efficiency | No sleep problems           | 69               | 34                   | 62            | 31                   |
|                             | Medium sleep problems       | 14               | 7                    | 17            | 9                    |
|                             | Serious sleep problems      | 11               | 5.2                  | 7             | 4                    |
|                             | Very serious sleep problems | 103              | 51                   | 113           | 57                   |
| 6 Sleep disturbances        | No sleep problems           | 22               | 11                   | 6             | 3                    |
|                             | Medium sleep problems       | 153              | 76                   | 171           | 86                   |
|                             | Serious sleep problems      | 26               | 13                   | 20            | 10                   |
|                             | Very serious sleep problems | 0                | 0                    | 0             | 0                    |
| 7 Use of sleep medications  | No sleep problems           | 172              | 86                   | 173           | 87                   |
|                             | Medium sleep problems       | 18               | 9                    | 14            | 7                    |
|                             | Serious sleep problems      | 9                | 5                    | 3             | 2                    |
|                             | Very serious sleep problems | 1                | 1                    | 7             | 1                    |
| 8 Daytime dysfunction       | No sleep problems           | 78               | 39                   | 52            | 26                   |
|                             | Medium sleep problems       | 66               | 33                   | 72            | 36                   |
|                             | Serious sleep problems      | 43               | 21                   | 58            | 29                   |
|                             | Very serious sleep problems | 14               | 7                    | 15            | 8                    |

As seen in this table, the majority of the students, dormitory or non-dormitory, have weak overall sleep quality and are mostly (80%) challenged on average or seriously in 7 dimensions of sleep. More than half of the students have serious sleep problem regarding habitual sleep efficiency. In order to evaluate the relation between residency status and sleep quality, statistical T test was performed the results of which are included in table 2.

As the results of table 2 suggest, there is significant relation between overall sleep quality and daytime dysfunction, and residence status ( $p < 0.05$ ), while there

is no significant relation between other sleep quality dimensions and residence status. Relation between sleep quality and its dimensions with educational status of the students are included in table 3.

As the results of table 3 suggest, the only dimension with significant relation with educational status of the students is the subjective sleep quality ( $p < 0.05$ ). In order to evaluate the residence status and educational status, independent statistical t test is used, the results of which is included in table 4.

As this table suggests, there is significant relation between the two mentioned terms ( $p < 0.05$ ).

**Table 2.** relation of sleep quality dimensions with residence status

| Sleep quality dimension     | Residence status |                    |               |                    | P-value* |
|-----------------------------|------------------|--------------------|---------------|--------------------|----------|
|                             | Dormitory        |                    | Non-dormitory |                    |          |
|                             | Average          | Standard deviation | Average       | Standard deviation |          |
| 1 Overall sleep status      | 7.36             | 2.56               | 7.97          | 2.55               | 0.018    |
| 2 Subjective sleep quality  | 1.09             | 0.75               | 1.18          | 0.79               | 0.254    |
| 3 Sleep latency             | 1.30             | 0.88               | 1.40          | 0.84               | 0.228    |
| 4 Sleep duration            | 1.03             | 1.07               | 1.09          | 1.08               | 0.576    |
| 5 Habitual sleep efficiency | 1.75             | 1.39               | 1.86          | 1.37               | 0.394    |
| 6 Sleep disturbances        | 1.02             | 0.49               | 1.07          | 0.36               | 0.366    |
| 7 Use of sleep medications  | 0.21             | 0.56               | 0.21          | 0.64               | 0.387    |
| 8 Daytime dysfunction       | 0.97             | 0.94               | 1.18          | 0.91               | 0.020    |

Significance level= 0.05

**Table 3.** relation of sleep quality and its sub-dimensions with educational status

| Sleep quality dimension     | Educational status |          |
|-----------------------------|--------------------|----------|
|                             | R2                 | P-value* |
| 1 Overall sleep status      | -0.061             | (0.232)  |
| 2 Subjective sleep quality  | -0.111             | (0.028)  |
| 3 Sleep latency             | 0.017              | (0.734)  |
| 4 Sleep duration            | -0.001             | (0.977)  |
| 5 Habitual sleep efficiency | -0.001             | (0.980)  |
| 6 Sleep disturbances        | 0.006              | (0.899)  |
| 7 Use of sleep medications  | 0.012              | (0.807)  |
| 8 Daytime dysfunction       | -0.093             | (0.064)  |

Significance level= 0.05

**Table 4.** relation of educational status with residence status

| Index              | Residence status |                    |               |                    | P-value* |
|--------------------|------------------|--------------------|---------------|--------------------|----------|
|                    | Dormitory        |                    | Non-dormitory |                    |          |
|                    | Average          | Standard deviation | Average       | Standard deviation |          |
| Educational status | -1.23            | 2.16               | 0.22          | 2.18               | 0.000    |

Significance level= 0.05

## Discussion

Results of this study suggest that educational status of non-dormitory students is better than the dormitory

ones. There is a set of factors causing educational failure that in turn, leads to loss in human and non-human resources. Therefore, different factors including:

motivation, objective, planning, facilities, individual and family conditions, economic and cultural status of the family and etc. must be considered in finding the causes of educational failure.<sup>15</sup> Residence in dormitory comes with exclusive conditions for students which can cause both educational failure and educational improvement. It can bring one into building relationship with a vast spectrum of students of different majors and grades. Comparison of job market of one's major with of the others can result in decrement of motivation and purpose and in turn, educational failure.

The researches performed, smoking by unsuccessful students is reported more than the successful ones with significant statistical difference.<sup>15</sup> This issue can stem from wasting time and spending spare time of students with friends in dormitories or outside smoking which naturally obstructs educational improvement. Since the dormitory students are in contact with many of their own age, they are more exposed to wasting time with playing computer games or other hobbies. According to interview of the researcher with some studied students, it was discovered that they were busy playing an online computer game or similar ones sometimes for several days and were not even present in the classes. Therefore, educational failure for dormitory students in comparison to their high school time when they were under control and support of the family is not unexpected.

Study results suggest that sleep quality mark for most of the students, dormitory or non-dormitory, was higher than 5 which means weak sleep quality. Qhoreishy et al studied sleep quality in students of Zanjan University of Medical Sciences in 1386, and come to conclusion that sleep quality which is an impactful factor for learning for them is low.<sup>16</sup> Study results suggest that the only dimension significantly related to educational status of the students is the subjective sleep quality. Farhady-Nasab et al concluded in a study named "study of subjective sleep quality and pattern in Medicine students of the Hamedan

University of Medical Sciences and its relation with characteristic adjectives that a significant number of medicine students suffer from sleep disturbances in night time sleep; a matter which can negatively affect health and daily performance.<sup>17</sup> Aside from the mentioned dimension, the others do not affect the sleep quality.

Ghavam et al concluded from a study performed in 1382-1383 named impact of occupation of dentistry students on their educational success in faculty of dentistry of the Iran University of Medical Sciences that occupation is a factor negatively affecting the educational status of the students<sup>18</sup> – a finding that complies with results of the present study. There is a set of factors causing educational failure including: motivation, objective, planning, facilities, individual and family conditions, economic and cultural status of the family and etc.<sup>15</sup> Considering that educational status is a complex category and is affected by different factors, it can be said that sleep quality can be under influence of other factors with lesser impact on educational status including the objective.

### Conclusion

There is a set of factors causing educational failure including: motivation, objective, planning, facilities, individual and family conditions, economic and cultural status of the family and etc. Considering that educational status is a complex category and is affected by different factors, it can be said that sleep quality can be under influence of other factors with lesser impact on educational status including the objective.

### Acknowledgement

The authors of the article are grateful to all the students who contributed to this study (Code of Ethics: IR.IUMS.REC 1395.95-09-153-29382).

### References

1. Purani H, Friedrichsen S, Allen AM. Sleep quality in cigarette smokers: Associations with smoking-related outcomes and exercise. *Addictive behaviors*. 2019;90:71-6.

2. Valiensi SM, Belardo MA, Pilnik S, Izbizky G, Starvaggi AP, Branco CC. Sleep quality and related factors in postmenopausal women. *Maturitas*. 2019;123:73-7.
3. Foulkes L, McMillan D, Gregory AM. A bad night's sleep on campus: an interview study of first-year university students with poor sleep quality. *Sleep Health*. 2019;5(3):280-7.
4. Ohlmann B, Bömicke W, Habibi Y, Rammelsberg P, Schmitter M. Are there associations between sleep bruxism, chronic stress, and sleep quality? *Dentistry*. 2018;74:101-6.
5. LeBlanc M, Beaulieu-Bonneau S, Mérette C, Savard J, Ivers H, Morin CM. Psychological and health-related quality of life factors associated with insomnia in a population-based sample. *Psychosomatic research*. 2007;63(2):157-66.
6. e Silva JAC .Sleep disorders in psychiatry. *Metabolism*. 2006;55:S40-S4.
7. Lima PF, Medeiros A, Araujo JF. Sleep-wake pattern of medical students: early versus late class starting time. *Brazilian journal of medical and biological research*. 2002;35(11):1373-7.
8. Johns MW, Dudley HAF, Masterton JP. The sleep habits, personality and academic performance of medical students. *Medical education*. 1976;10(3):158-62.
9. Hidalgo M, Caumo W. Sleep disturbances associated with minor psychiatric disorders in medical students. *Neurological sciences*. 2002;23(1):35-9.
10. Veldi M, Aluoja A, Vasar V. Sleep quality and more common sleep-related problems in medical students. *Sleep medicine*. 2005;6(3):269-75.
11. Dahlin ME, Runeson BO. Burnout and psychiatric morbidity among medical students entering clinical training: a three year prospective questionnaire and interview-based study. *BMC Medical education*. 2007;7(1):6.
12. Tyssen R, Røvik JO, Vaglum P, Grønvold NT, Ekeberg Ø. Help-seeking for mental health problems among young physicians: is it the most ill that seeks help? *Social psychiatry and psychiatric epidemiology*. 2004;39(12):989-93.
13. Buysse DJ, Reynolds III CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry research*. 1989;28(2):193-213.
14. Heydari AR, Ehtesham Zadeh P, Marashi M. Relationship between the severity of insomnia, sleep quality, sleepiness and mental health disorder with academic performance in girls. *Woman and Culture*. 2010;1(4):65-76. [Persian]
15. Rahimi PS, Heshmati H, Moghaddam Z. Survey the frequency and risk factors in the academic failure in students that living in dorms of golestan university of medical sciences & health sciences in year 2011-2012. 2014;2(1):16-27.
16. Ghoreishi A, Aghajani AH. Sleep quality in Zanjan university medical students. *Tehran university medical journal TUMS publications*. 2008;66(1):61-7. [Persian]
17. Farhadi Nasab A, Azimi H. Study of patterns and subjective quality of sleep and their correlation with personality traits among medical students of Hamadan University of Medical Sciences. *Avicenna journal of clinical medicine*. 2008;15(1):11-5.[Persian]
18. Ghavam M, Pourmalek F, Imani V. Effect of densitry students' employment on their academic success. (2003-2004). *Islamic dental association of iran (majallah-i-dandanpizishki)* 2005;17(1):104-12.