

Association between Maximum Aerobic Power and General Health in Women

Ehsanollah Habibi¹, Moslem Mombeini^{2*}, Zahra Ordudari³, Hadis Noorouzi⁴, Ghasem Yadegarfar⁵, Zahra Koolivand⁶

¹ Professor, Department of Occupational Health Engineering, School of Health, Isfahan University of Medical Sciences, Isfahan, Iran² MSc Student, Department of Occupational Health Engineering, School of Health, Isfahan University of Medical Sciences, Isfahan, Iran³ MSc Student, Department of Occupational Health Engineering, School of Health, Isfahan University of Medical Sciences, Isfahan, Iran⁴ MSc Student, Department of HSE Management, Islamik Azad University Najafabad Branch, Isfahan, Iran⁵ Assistant Professor, Biostatistics Department, School of Health, Isfahan University of Medical Sciences, Isfahan, Iran⁶ Occupational Health Expert, Department of Occupational Health Engineering, School of Health, Hamedan University of Medical Sciences, Hamedan, Iran^{*} Corresponding author: Moslem Mombeini, Email: mosimomsal@gmail.com, Tel: +98-913-7712947

Abstract

Background: The highest aerobic capacity (VO_{2max}) is one of the criteria that can be used to create a balance between work and worker. To maintain the health and physical capacity of individuals, as well as to increase production and profitability, physical work capacity (PWC) can be used to put individuals in jobs proportional to their capacities.

Methods: This cross-sectional study was performed on 120 women working in hospitals aged 21-49 years old. Individuals were evaluated by a single-stage Ebbeling test. A questionnaire about general health, job fatigue and job satisfaction was used, along with a treadmill. Data were analyzed by SPSS V22 software using the Pearson correlation and t-test.

Results: According to the results, the highest aerobic capacity, physical working capacity and general health were 38·89(1·85) ml/kg body weight per minute, 4·10(0·47) kcal/min and 25·43(12·09) respectively, and mean and standard deviation of fatigue and job satisfaction scores were 85·87(32·35) and 57·47(13·01) respectively. In this study, there was a significant relationship between VO_{2max} and age, weight, body mass index (BMI), fatigue and job satisfaction. **Conclusion:** According to the results, since fatigue and job satisfaction are related to workers' VO_{2max}, it is recommended that, in addition to considering the aspects of physical fitness in the workplace, special attention should be paid to mental aspects, since they affect the physical and, consequently, the productivity of workers.

Keywords: Maximum aerobic capacity; Physical work capacity; Hospital women employees; General health

Introduction

The main objective of ergonomic science is to create a balance between people and their environment. For this reason, tools for assessing the effectiveness of ergonomic interventions are needed to provide the necessary information on proportionality vs. disproportionality.¹ To provide such

a ratio, it is necessary to determine the energy consumption of different occupations and, on the other hand, to measure the physical capacity of the employees. Therefore, measuring physical fitness is one of the key issues of ergonomic science, which has been emphasized by the International Labor Organization.²

Citation: Habibi E, Mombeini M, Ordudari Z, Noorouzi H, Yadegarfar Gh, Koolivand Z. **Volatile Organic Compounds Concentration in the Laboratory Medical Laboratories.** Archives of Occupational Health. 2018; 2(3): 148-55.

Article History: Received: 4 March 2018; Revised: 22 April 2018; Accepted: 1 July 2018

Copyright: ©2017 The Author(s); Published by Shahid Sadoughi University of Medical Sciences. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In other words, by measuring the mental, physical and physiological characteristics of humans, all related factors can be tailored to fit the work, along with his measurements and physiological capacities.³ In Iran, there is no accurate statistics on the employment rate of the workforce in the treatment sector. In Europe, however, women account for the highest percentage of civilian personnel. As much as 75% of these forces are women. in addition, the distinction between work in the hospital and the production industries is inevitable.⁴

Hospital workers are required to work round the clock.⁵ Working in the field of treatment, and, in particular, nursing, is one of the areas where physical and psychological stimuli (overtime, difficulty and excessive workload, unwanted overtime) threaten comfort and quality of life.⁶ The disparity between human abilities and the type of work that they do causes problems and issues.⁷ One of the important aspects of ergonomics and organizational management is work physiology,^{8,9} which involves the selection of the required workforce and the work-life balance of individuals with physical and mental capabilities. By measuring individuals physiologically, one can place each person in a career appropriate for his physiological threshold, in addition to maintaining his health. Further, physical capacity, production and productivity will also increase.³ In 1962, Professor B. Bink coined the term Physical Work Capacity (PWC), which refers to the maximum amount of energy that a person can perform during his eight-hour shift without sacrificing his own health.³

VO2-max is the highest amount of oxygen that can be absorbed by the respiratory tract and transmitted through the blood to the muscles, and is considered as a standard for measuring the practical limits of the cardio-respiratory system.^{3,10} VO2-max is usually expressed in liters or milliliters of oxygen per kg of body weight per minute (1 kg⁻¹ min⁻¹ or ml kg⁻¹ min⁻¹).¹¹ VO2-max varies between different individuals and is affected by various factors such as physical,

psychological, environmental and genetic.^{10,11,12} Aerobic capacity can be measured directly or indirectly. Direct assessment of aerobic capacity is commonly used in young and athletic subjects. Indirect measurement, which has a lower physiological pressure on a person, is more suitable for working communities.¹¹

From an economic standpoint, the correct assessment of employees' ability to work is important for the employer, and strengthening it is one of the ways to increase the productivity of human resources in industries and organizations. The ability to work is the basis of the design of work stations and the screening of potential employees for specific occupations. In other words, if the physical and psychological abilities of workers and employees are not in line with their occupational needs, it leads to safety and health problems, reduces production, and results in increased dismissal of workers and employees, which adds to expenditure.¹³

Despite numerous studies on male hospital and non-hospital workers, no study has been conducted to determine the relationship between VO2MAX physical factors and psychological factors (general health) in women working in hospitals. This study aimed to measure this relationship. If there is a direct relationship between mentioned factors, in addition to considering the physical aspects of balance of work, special attention should be paid to the psychological suitability of work with female personnel. The aim of this study was to determine the maximum aerobic power (VO2max), physical working capacity (PWC) and general health (GH) and to identify underlying factors such as age, weight, BMI, fatigue and job satisfaction in female employees working in Isfahan hospitals.

Methods

This cross-sectional descriptive-analytic study was carried out on 120 women working in Isfahan hospitals in 2014. The sample size needed for this study with a confidence level of 95% and a durability

interval of $d = 2$, and considering the possibility of data loss using the formula $N = z^2 * sd^2 / d^2$, was estimated to be 96. The sample, following the model of previous studies, was selected randomly from a population aged 20 to 49 years.^{6,14} To ensure the results achieved and to increase the credibility of the study, the sample size was increased to 120 people. It should be noted that the participants filled out an informed consent form before completing the tests and were assured that the collected data would be kept confidential and would not be provided to any entity. Also, those who suffered from cardiovascular problems or respiratory problems were excluded from the study. At the beginning of the test, a questionnaire for gathering information such as additional employment, exercise patterns, whether participants smoke, marital status, work system, type of work and level of education was given to each individual. In this study, a portion of the data was collected by a questionnaire and another part by a direct measurement, which is described below:

In this study, the GHQ-28 questionnaire was used to determine the general health of individuals. Validity and reliability of the questionnaire have been confirmed in several studies.¹⁵ The 28-item form of the General Health Questionnaire was developed by Goldberg DP and Hillier VF in 1979, which includes four subscales of physical symptoms, anxiety and sleep disorders, social dysfunction and severe depression.¹⁵ In these subscales, those who score 6 or less are considered healthy, while those who score 7 or higher are suspected of being impaired. With this delineation, the sensitivity and specificity of the test were 84.7(2.7) and 93.7%(2.7) respectively, and the overall classification error was 8.2 (2.7).¹⁶ Also, in this method, the score for those who score 23 or less in overall health are considered healthy and those who score 24 or more are suspected of having the disorder. Sensitivity, specificity and error rate of

general classification with this cutting score were 70.5 (2.4), 92.3 (2.4) and 12.3 (2.4) respectively.¹⁶

Brayfield AH and Rothe Job Satisfaction Questionnaire: The questionnaire was created by Brayfield AH, Rothe in 1951 to measure job satisfaction and includes 19 5-option questions. In this questionnaire, respondents are asked to identify their feelings and attitudes toward their jobs on a 5-point scale. For the options strongly disagree, disagree, no idea, agree and strongly agree, the score of 1, 2, 3, 4, and 5 are assigned, respectively. The rest of the items are scored in reverse. The reliability of the Brayfield AH and Rothe job satisfaction questionnaire has been confirmed in several studies in Iran.¹⁷

Swedish Occupational Fatigue Questionnaire (SOFI-20): The Persian version of the Swedish Job Fatigue Questionnaire has 20 items and 5 dimensions: "Lack of energy", "Physical effort", "Physical discomfort", "Lack of motivation" and "Sleepiness". Each item is rated using an 11-degree Likert scale from zero (completely disagree) up to 10 (completely agree). The minimum and maximum scores are 0 and 200 respectively. The reliability and validity of the questionnaire are confirmed in various studies.¹⁸

Method for measuring the maximum aerobic capacity (VO2-max)

In this study, the VO2-max was determined with the treadmill exercise test (using a T 9670 treadmill) and Ebbeling method.¹⁹

To determine the PWC, the obtained VO2max (in $l\ min^{-1}$) will be multiplied by 5 to achieve maximum physical ability (AC). Finally, the physical working capacity of each person was determined from the equation $PWC = AC \times 0.33$ in kilocalories per minute.²⁰

The height and weight of the subjects in the study were measured in light clothing, without shoes or gloves. For measuring heartrate and blood pressure, a polar sport tester and the blood pressure gauge device were used respectively. Body mass

index was also obtained by dividing the weight by height and in square kilograms.

After experiments, the collected data were analyzed by SPSS software version 22 using the t-test with two independent samples, Spearman's rank-order correlation, analysis of variance, and the correlation coefficient formula. It should be noted that the quantitative data from the mean and standard deviation with 95% confidence intervals were used, and for the qualitative data, 95% confidence intervals were used.

Results

The average age, work history, weight, height, and body mass index of the subjects are shown in Table

1. Also, the number and percentage of studied variables (general health, satisfaction, and occupational stress) has been shown in the participants in Table 2.

The relationship between VO₂max and age, weight, height, BMI and general health score is shown in Table 3. As shown in this table, with the increase in age, the maximum aerobic power in humans is reduced. In this table, based on the general health score and its sub-indices, the difference in mean of maximum aerobic capacity in healthy and suspected cases is statistically significant, so that the maximum aerobic capacity is higher in healthy people.

Table 1. Average demographic information, maximum aerobic capacity, physical activity capacity, general health score and sub-scales, fatigue and job satisfaction in the studied population.

Variable	mean	SD	Min	Max
Age in years	34.47	6.29	21	49
Height in meters	1.61	0.05	1.50	1.78
Weight in kilograms	64.06	8.22	42	88
BMI in kilograms square meters	24.52	2.95	17.58	33.53
Work experience	11.20	6.71	2	23
VO ₂ MAX in ml min ⁻¹ kg ⁻¹	38.89	1.85	33.07	45
VO ₂ MAX in l min ⁻¹	2.48	0.28	1.70	3.21
PWC in kcal min ⁻¹	4.10	0.47	2.80	5.30
Overall health score	25.43	12.09	6	70
Physical symptoms scale	8.20	3.98	00	17
Anxiety symptoms and sleep disorders	7.10	4.51	00	21
Social disorder scale	6.92	2.78	1	18
The scale of symptoms of depression	3.20	3.66	00	19
Heartbeat	127.30	8.04	109	151
Job fatigue	85.87	32.35	16	184
Job satisfaction	57.48	13.1	24	80

Table 2. The percentage and frequency of demographic information of the participants

Variable	Subgroup	Number	Percent
Second job	Yes	24	%20
	No	96	%80
Exercise	Yes	41	%34.2
	No	79	%65.8
Smoke	Yes	9	%7.5
	No	111	%92.5
Marital status	Single	24	%20
	Married	96	%80
Work system	Day work	66	%55
	Shift	54	%45
Kind of work	Official	45	%37.5
	Executive	75	%62.5
Education	Diploma and lower	23	%19.2
	Associate degree	14	%11.07
	Bachelor's degree and higher	83	%69.2

Table 3. Relationship between age, weight, height, BMI and general health score with VO2max

Variable	Group	Number	Percent	VO2max (ml kg-1 min-1)				
				Mean	SD	MIN	MAX	P-Value*
Age	20-29	37	%30.8	40.28	1.59	37.11	45	0.001
	30-39	47	%39.2	38.74	1.67	33.07	41.16	
	40-49	36	%30	37.66	1.34	33.92	39.87	
Weighted groups	≤55	22	%18.3	40.03	1.60	37.65	45	0.001
	56-64	41	%34.2	38.97	1.33	36.78	43.75	
	65-69	27	%22.5	39.01	1.73	34.42	41.97	
	≥70	30	%25	37.84	2.23	33.07	42.29	
Height groups in cm	≤155	18	%15	39.20	1.03	37.65	41.16	0.022
	156-160	38	%31.6	39.24	1.57	36.78	45	
	161-165	32	%26.7	38.02	2.35	33.07	43.75	
	≤166	32	%26.7	39.17	1.76	34.11	42.29	
BMI groups	18.5<	1	%0.8	39.67	0	39.67	39.67	0.001
	18.5-24.9	70	%58.3	39.28	1.75	34.42	45	
	25-29.99	40	%33.3	38.71	1.55	33.92	41.41	
General health score	≥30	9	%7.5	36.62	2.36	33.07	41.16	0.001
	0-23	64	%53.3	39.73	1.60	36.49	45	
	24-84	56	%46.7	37.93	1.66	33.07	40.69	
Physical symptoms scale	0-6	51	%42.5	39.57	1.86	34.11	45	0.001
	7-21	69	%57.5	38.39	1.69	33.07	41.94	
Sleep	0-6	65	%54.2	39.52	1.59	36.49	45	0.001
	7-21	55	%45.8	38.15	1.88	33.07	41.94	
Disruption of social function	0-6	49	%40.8	39.63	1.54	36.49	45	0.001
	7-21	71	%59.2	38.38	1.89	33.07	43.75	
The scale of symptoms of depression	0-6	100	%83.3	39.15	1.68	33.92	45	0.001
	7-21	20	%16.7	37.59	2.14	33.07	40.46	

* statistical test: t-Test** p<0.05

Table 4. Relationship between heart rate, fatigue and job satisfaction with VO2max

Variable	Mean	SD	MIN	MAX	VO2max (ml kg-1 min-1)		P-Value*
					R Correlation Coefficient	P-Value*	
Heartbeat	127.30	8.04	109	151	-0.034	0.714	
Job fatigue	85.87	32.35	16	184	-0.335	0.001	
Job satisfaction	57.48	13.01	24	80	0.331	0.001	

*p<0.05

As shown in Table 4, there is a negative correlation between job fatigue and VO2max and decreases with increasing this VO2max variable, but there is a positive correlation between job satisfaction and VO2max. As a result, there is a significant relationship between job satisfaction and occupational exhaustion with VO2max. With the increase in occupational fatigue, the maximum aerobic capacity is reduced, as well as by increasing job satisfaction, the maximum aerobic capacity increases. Meanwhile, there was no significant relationship between heartrate and maximum aerobic capacity in this study. P = 0.714.

In Table 5, the relationship between second job, exercise, whether participants smoked, marital status, work system and type of work with VO2max were investigated. As can be seen in this table, there is a statistically significant relationship between the two variables of career, marriage, and work system with aerobic capacity. Meanwhile, aerobic capacity is higher among people in one job. Also, in day workers and single people, aerobic capacity is higher than that of married and shift workers. In this study, there was no relationship between education level, type of work and smoking history with aerobic capacity.

Table 5. The relationship between the second job, exercise, smoking, marriage, work system and type of work with VO2max

Variable	Subgroup	VO2max (ml kg-1 min-1)	Mean	SD	P-Value*
Second job	Yes	37.78	1.76	0.001	
	No	39.17	1.78		
Exercise	Yes	39.68	1.82	0.001	
	No	38.49	1.75		
Smoke	Yes	38.90	1.12	0.989	
	No	38.89	1.90		
Marital status	Single	40.17	1.57	0.001	
	Married	38.57	1.79		
Work system	day work	39.27	2.01	0.013	
	shift	38.43	1.53		
Type of work	Official	38.72	1.31	0.429	
	Executive	39	2.12		

p<0.05*

Discussion

In the present study, the mean age of the studied population was 34.47 years and most of the subjects were in the natural range of BMI. Mean VO2max and PWC were 38.89 ml kg-1 body weight per minute and 4.10 kcal min-1, respectively. According to Bugajskaa et al., the mean VO2max values were 9.8 (6.33), which is slightly different from the current study.²¹ Also, the mean of total general health, fatigue and job satisfaction in the study population was **25.43(12.09)**, **85.87(32.35)**, and **57.47(13.01)**. Based on the results of the general health score, it can be judged that the ratio of healthy people with a low percentage is more than those who are suspected of mental illness. The results of this study showed that among the subscales of general health, the mean scores below the scale of physical symptoms, sleep, and anxiety and disruption of social function in the subjects exceeded the delineation (6 points), and only the mean score and only the sub-scales of depression symptoms are lower than the delineation.

The results of this study are consistent with the findings of the study by Kessler et al. in the two sub-comparisons of physical symptoms and anxiety.²² This study is less consistent, however, with the results of Stanfield et al. who show in their study even though they have less work experience, fewer

psychiatric disorders are observed in women.¹⁵ In this study, as well as the results of the research by Fakhar et al., there is a statistically significant relationship between psychiatric disorders in married and single people, and married people are more likely to experience mental disorders.¹⁴ In the study of Bugajskaa et al., it was observed that aerobic capacity decrease with the increase of age, which is consistent with the results of the present study.²¹

In this study, the mean values and standard deviations of VO2-max in ml kg-1 min-1 were higher than those obtained in the Bugajskaa et al. study.²¹ Considering the calculation of physical working capacity (PWC) by the highest aerobic capacity, it can be concluded that the physical working capacity in Kcal min-1 obtained in this study is the same as the mean of the maximum aerobic capacity of the mean obtained in the study of Bugajskaa et al.²¹ In contrast to the research of Moghiseh et al., no significant correlation was found between the maximum aerobic capacity of VO2-max and heart rate in this study.²³ The results of this study showed that increasing the work record and BMI did not increase the overall health score. These findings are not consistent with the results of other studies.^{16,24} Also, according to the results of this study, there is a significant relationship between the aerobic capacity of individuals with job satisfaction and the feeling of fatigue during work, which is contrary to the findings of other studies.^{1,25} These differences seem to be due to differences in the statistical society.

Conclusion

This study showed that there is a significant statistical relationship between mental health and maximum aerobic capacity. The results of this study confirmed the findings of previous studies on the association of maximum aerobic capacity with weight, height, age, and body mass index. Also, in this study, job fatigue and job satisfaction are among the factors affecting general health and

aerobic capacity. It seems that mental health in women working in hospitals requires special consideration. In this area quantitative studies are needed regarding the maximum aerobic capacity, physical capacity, and general health and parameters affecting them, and it is recommended that more research be done by the researchers. Also, in addition to considering the physical aspects, psychological appropriateness of work with the personnel also should be specially taken into consideration.

Conflict of interest

The authors declare no conflict of interests

Acknowledgment

The authors of the article express their appreciation and thanks to the authorities and all medical personnel of hospitals in Isfahan. This study is based on Master's thesis of Mr. Moslem Mombeini, student of Occupational Health Engineering, Isfahan University of Medical Sciences with research code 394737.

References

- Choobineh AR, Barzideh M, Gholami T, Amiri R, Tabatabaei HR, Almasi HA. Estimation of Aerobic Capacity (VO₂-max) and Study of Its Associated Factors among Male Workers of Industrial Factories in Sepidan/Fars Province. *Archive of SID*, 2009. 10(1):1-12 In Press. [Persian]
- Motalebi Kashani M, Lahmi MA, Khavanin A. Evaluation of physical work capacity at Iranian workers. *ICE07 Kuala Lumpur*. 2007;42-47. [Persian]
- Mououdi MA, Choobineh AR. Ergonomics in practice: selected ergonomics topics. Tehran: Nashr-e-Markaz. 1999;81-94. [Persian]
- Ali Arabian F, Motamedzade M, golmohammadi R, Moghim Beigi A, Pir Hayati F. The impact of ergonomics intervention on musculoskeletal disorders among nahavand alimoradian hospital staff. *Ergonomics*. 2013;1(1):23-32. [Persian]
- Arghami Sh, Ghanbari H, Zahirian Gh. Health status of city staff shiftwork in a hospital Razan. *Ergonomics*. 2014;2(1):64-69. [Persian]
- Feyzi V, Jafari Roodbandi AS, Farahbakhsh S, Rezaei H. The investigation of occupational and demographic factors effective on the quality of life of nurses and nurse aides working in teaching hospitals affiliated to Kerman university of medical sciences in 2014. *Ergonomics*. 2016;4(3):33-40. [Persian]
- Motalebi Kashani M, Lahmi MA, Khavanin A. Evaluation of physical work capacity at Iranian workers. *ICE07 Kuala Lumpur*. 2007;42-47. [Persian]
- Wong DP, Carling C, Chaouachi A, Dellal A, Castagna C, Chamari K, et al. Estimation of oxygen uptake from heart rate and ratings of perceived exertion in young soccer players. *Strength & conditioning research*. 2011;25(7):1983-8.
- Motalebi M. Assessment of work ability in workers employed in machining industry and provide appropriate scientific methods for selecting workers for physical work, [Doctorate Thesis]. Iran. Teacher Training University;1994. [Persian]
- Uth N, Sørensen H, Overgaard K, Pedersen PK. Estimation of VO₂max from the ratio between HR_{max} and HR_{rest} – the heart rate ratio method. *European journal of applied physiology*. 2004;91(1):111-5.
- Tayyari, F. Smith, J L. "Occupational Ergonomics" Chapman & Hall, first edition, U.S.A. PP. 108-113. 1997.
- Valipour F, Khavanin A, Asiliyan H, Pourtaghi Gh. H, Mohebi HA, Jonaidi N, et al. Measurement of physical work capacity (pwc) for iranian military personnel in different condition chamber laboratory clime (Normal and Very Heat Humid). *Journal of Military Medicine* 2007; 9(1): 67-72. [Persian]
- Habibi E, Dehghan H, Zeinodini M, Yousefi H A, Hasanzadeh A. The Relation between Work Ability Index and Physical Work Capacity Based on Fox Equation for VO₂ Max in Male Nursing Staff of Isfahan Hospitals, Iran. *Journal of Health System Research*. 2012 jan and feb; 7(5): 606-614. [Persian]
- Fakhari, A. Ranjbar, F. Dadashzadeh, H. Moghaddas, F. An epidemiological survey of mental disorders among adults in north, west area of Tabriz, Iran. *Pakistan J of Med*. pp. 23:N1. [Persian]
- Stanfeld, S.A. Marmot, M.G. Social class and minor psychiatric disorder in British Civil Servants: a validated screening survey using the General Health Questionnaire. *Psychol Med*. 22, pp. 39-49.
- Noorbala AA, Bagheri yazdi SA, Mohammad K. The Validation of General Health Questionnaire- 28 as a Psychiatric Screening Tool. *Hakim Research Journal*. 2009;11:47- 53. [Persian]
- Rezaei A, Khalil zadeh A. The relationship between social intelligence and job satisfaction of school administrators. *JOURNAL OF EDUCATIONAL SCIENCES*, 2009; 2(7): 121-145. [Persian]
- Javadvour F, Keshavarzi S, Choobineh A, Aghabaigi M. Validity and reliability of the Swedish Occupational Fatigue Inventory (SOFI-20) among Iranian working population. *Journal of Ergonomics*, Spring 2015; 3(1): 50-58. [Persian]
- Ebbeling CB, Ward A, Puleo EM, Widrick J, Rippe JM. "Development of a single-stage submaximal treadmill walking test". *PubMed - indexed for MEDLINE*. See comment in PubMed Commons below1991 Aug; 23(8): 966-73.
- Hosseiniabadi S, Hamidi Bagejan P, Ebrahimi H, Barkhordari A, Raie Bandepe T. Estimation of Aerobic Capacity (VO₂-max) and Physical Work Capacity in Laborers. *Knowledge & Health* 2013;8(3):131-137. [Persian]
- Bugajska J , Makowiec-Dąbrowska T, Jegier A , Marszalek A. Physical work capacity (VO₂ max) and work ability (WAI) of active employees (men and women) in Poland. *International Congress Series* 1280 (2005) ; 156–160.

22. Kessler, R.C. Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United State. 51, pp. 8-19.
23. Moghiseh M, Habibi E, Aramesh N, Hasanzadeh A, Khorvash MK, Poorrahmatian AH. The association between VO₂max and heart rate of casting industry workers. JOHE, Winter-spring 2013; 2 (1-2): 20-26.
24. Choobineh AR, Neghab M, Hasanzadeh J, Rostami R. Comparative Assessme t of e tists' Psychological Health Status in Shiraz with Their Physicians Counterparts Using General Health Questionnaire (GHQ-28). Journal of Dental Medicine-Tehran University of Medical Sciences 2013; 25:290-96. [Persian]
25. Virtanen M, Vahtera J, Pentti J, Honkonen T, Elovainio M, Kivimki M. Job strain and psychologic distress influence on sickness absence among Finnish employees. Am J Prev Med 2007;33(3):182-7.